



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN STATISTICAL REPORTS.

ARABIA—*Aden*.—Month of April, 1899. Estimated population, 30,000. Total number of deaths, 104, including 50 from enteric fever.

Month of May. Total number of deaths, 109, including smallpox, 1, and 48 from enteric fever.

BAHAMAS—*Dunmore Town*.—Two weeks ended August 4, 1899. Census population, 1,472. No deaths and no contagious diseases reported.

Governors Harbor.—Week ended August 5, 1899. Estimated population, 1,500. No deaths and no contagious diseases reported.

Green Turtle Cay—Abaco.—Two weeks ended August 3, 1899. Estimated population, 3,900. No deaths and no contagious diseases reported.

Inagua.—Four weeks ended July 30, 1899. Estimated population, 1,200. No deaths and no contagious diseases.

Nassau.—Two weeks ended August 5, 1899. Estimated population, 12,000. No deaths and no contagious diseases reported.

BRITISH HONDURAS—*Belize*.—Month of June, 1899. Estimated population, 30,000. Total number of deaths, 68, including 5 from phthisis pulmonalis.

DUTCH GUIANA—*Paramaribo*.—Month of June, 1899. Estimated population, 30,567. Total number of deaths, 80. No contagious diseases.

GERMANY—*Dresden*.—Month of June, 1899. Estimated population, 395,300. Total number of deaths, 592, including diphtheria, 3; enteric fever, 1; measles, 1; whooping cough, 4, and 86 from phthisis pulmonalis.

Glauchau.—Month of June, 1899. Estimated population, 27,000. Total number of deaths, 69, including diphtheria, 1; measles, 2; whooping cough, 3, and 9 from phthisis pulmonalis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended July 29, 1899, correspond to an annual rate of 20.7 a thousand of the aggregate population, which is estimated at 11,404,408. The highest rate was recorded in Liverpool, viz, 29.7, and the lowest in Wolverhampton, viz, 12.9.

London.—One thousand eight hundred and eighty-four deaths were registered during the week, including measles, 56; scarlet fever, 5; diphtheria, 35; whooping cough, 19; enteric fever, 13, and diarrhea and dysentery, 350. The deaths from all causes correspond to an annual rate of 21.6 a thousand. In Greater London 2,481 deaths were registered, corresponding to an annual rate of 19.8 a thousand of the population. In the "outer ring" the deaths included 9 from diphtheria, 11 from measles, and 11 from whooping cough.

Sunderland.—Two weeks ended July 22, 1899. Estimated population, 145,613. Total number of deaths, 80, including measles, 2, and 4 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended July 29, 1899, in the 22 principal town districts of Ireland was 22.0 a thousand of the population, which is estimated at 1,053,188. The lowest rate was recorded in Lurgan, viz, 21.6, and the highest in Sligo, viz, 40.6 a thousand. In Dublin and suburbs 191 deaths were registered, including diphtheria, 2; enteric fever, 11; measles, 13; scarlet fever, 3, and whooping cough, 3.

Scotland.—The deaths registered in 8 principal towns during the week ended July 29, 1899, correspond to an annual rate of 17.1 a thousand of the population, which is estimated at 1,587,414. The lowest mortality was recorded in Leith, viz, 11.4, and the highest in Perth, viz, 20.3 a thousand. The aggregate number of deaths registered from all causes was 523, including diphtheria, 3; measles, 17; scarlet fever, 6, and whooping cough, 9.

JAMAICA—Kingston.—Month of July, 1899. Estimated population, 46,542. Total number of deaths, 126, including 9 from phthisis pulmonalis.

JAPAN—Formosa—Tamsui.—Month of June, 1899. Estimated population, 2,797,543. Number of deaths not reported. Deaths from plague, 187.

Two weeks ended July 12. Total number of deaths not reported. Deaths from plague, 46.

MALTA.—Two weeks ended June 30, 1899. Estimated population, 180,328. Total number of deaths, 203, including enteric fever, 2; measles, 1, and 1 from whooping cough.

MEXICO—Tuxpan.—Month of July, 1899. Estimated population, 11,000. Total number of deaths, 41, including yellow fever, 1; measles, 1; whooping cough, 1; leprosy, 1, and 2 from phthisis pulmonalis.

PARAGUAY—Asuncion.—Estimated population, 45,000. Total number of deaths, 53, including enteric fever, 2, and 6 from phthisis pulmonalis.

RUSSIA—Riga.—Month of March, 1899. Estimated population, 282,000. Total number of deaths, 574, including diphtheria, 13; enteric fever, 5; measles, 4; scarlet fever, 3; whooping cough, 6; smallpox, 35, and 76 from phthisis pulmonalis.

Month of April. Total number of deaths, 537, including diphtheria, 10; enteric fever, 3; measles, 3; scarlet fever, 4; whooping cough, 3; smallpox, 34, and 66 from phthisis pulmonalis.

WEST INDIES—St. Thomas.—Months of April, May, and June, 1899. Estimated population, 14,007. Total number of deaths, 288.